

IBUPROFEN 200MG TABLETS

PATIENT INFORMATION LEAFLET

Read this leaflet carefully because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor or pharmacist if your symptoms worsen or do not improve.

IN THIS LEAFLET:

- 1. What this medicine is for**
- 2. Before you take the medicine**
- 3. How to take the medicine**
- 4. Possible side effects**
- 5. Storing the medicine**
- 6. Further information**

1. WHAT THIS MEDICINE IS FOR

Ibuprofen 200mg Tablets contain ibuprofen, which belongs to a group of medicines called Non-Steroidal Anti-Inflammatory Drugs (NSAIDs). These medicines work by changing how the body responds to pain, swelling and high temperature.

This medicine is used to relieve:

- rheumatic and muscular pain, backache
- neuralgia (nerve pain)
- headache, migraine
- dental pain, period pain
- fever (high temperature)
- the symptoms of cold and flu.

2. BEFORE YOU TAKE THE MEDICINE

Do not take Ibuprofen 200mg Tablets if you:

- have or have had a stomach ulcer, perforation or bleeding of the stomach
- are allergic to ibuprofen, to any of the ingredients (listed in section 6), or to aspirin or other painkillers (an allergic reaction may be recognised as shortness of breath, runny nose, skin rash or itching)
- have kidney or heart failure or severe liver failure
- are taking more than 75mg of aspirin a day
- are in the last 3 months of pregnancy.

Talk to your doctor or pharmacist before taking ibuprofen tablets if you:

- are elderly because you may be at more risk of having serious side effects, particularly stomach problems
- have or have suffered from asthma, diabetes, high cholesterol or have allergies
- have liver or kidney problems
- have stomach or bowel disorders including Crohn's disease or a condition known as ulcerative colitis
- have Systemic Lupus Erythematosus (SLE) – an illness which affects your immune system. It causes joint pains, skin changes and problems with other parts of your body.
- have, have previously had, or are at risk of heart problems, high blood pressure or stroke. Medicines such as Ibuprofen 200mg Tablets may be associated with a small increased risk of heart attack (myocardial infarction) or stroke. Any risk is more likely with high doses and prolonged treatment. Do not exceed the recommended dose or duration of treatment (10 days). If you have heart problems, previous stroke or think that you might be at risk of these conditions (for example if you have high blood pressure, diabetes or high cholesterol or are a smoker) you should discuss your treatment with your doctor or pharmacist.

If you are taking other medicines

You must not take these tablets if you are taking certain other medicines – see section 2 'Do not take Ibuprofen 200mg Tablets if'.

Talk to your pharmacist or doctor before you take the tablets if you are taking any regular medication, especially:

- low –dose aspirin (up to 75mg daily)
- other medicines for thinning the blood (anti-coagulants (against clotting) and anti-platelet agents such as warfarin and ticlopidin)
- medicines for high blood pressure (ACE-inhibitors such as captopril, beta-blockers and angiotensin II antagonists) and water tablets (diuretics)
- corticosteroids
- methotrexate (a medicine for cancer)
- cardiac glycosides (medicines used to treat heart failure)
- cyclosporin and tacrolimus (immunosuppressant medicines often used following organ transplants)
- mifepristone (a medicine used to terminate pregnancy – NSAIDs should not be used for 12 days after mifepristone)
- lithium (for depression or mental problems)
- zidovudine (a medicine to treat viruses)
- quinolone antibiotics (medicines used to treat bacterial infections).

Pregnancy and Breastfeeding

Do not take ibuprofen tablets if you are in the last 3 months of pregnancy. Talk to your doctor before taking ibuprofen tablets if you are in the first 6 months of pregnancy or are breastfeeding.

Ibuprofen 200mg Tablets belong to a group of medicines which may affect fertility in women. Fertility goes back to normal when you stop taking the medicine. It is unlikely that if you only take

these tablets occasionally it will affect your chances of becoming pregnant. If you have problems becoming pregnant talk to your doctor before taking this medicine.

Important information about some of the ingredients of this medicine.

This product contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

3. HOW TO TAKE THIS MEDICINE

Age	Dose and how often to take
Adults, the elderly and children over 12 years	Take 1 to 2 tablets up to 3 times a day, as required. Swallow tablets with a glass of water preferably with or after food. Do not take more often than every 4 hours. Do not exceed 6 tablets in 24 hours.
<ul style="list-style-type: none">• This medicine is for short-term use only• Take the lowest dose for the shortest time necessary• Do not take Ibuprofen 200mg Tablets for longer than 10 days If you do not get better, or get worse, talk to your doctor. They will tell if it is safe to carry on taking this medicine.	
Do not give to children under 12 years.	

If you take to many tablets: Talk to a doctor straight away, or go to your nearest hospital casualty department. Take the carton and this leaflet with you.

4. POSSIBLE SIDE EFFECTS

Most people take Ibuprofen 200mg Tablets without any problems, but it can have side effects, like all medicines.

To reduce the chance of side effects, especially if you are elderly, use the lowest effective dose for the shortest possible time.

If you get any of the following at any time during your treatment STOP TAKING, and get medical help straight away:

Allergic reactions:

The following reactions may mean you are having an allergic reaction to this medicine:

- asthma, worsening of asthma, unexplained wheezing or shortness of breath
- swelling of the face, lips, tongue and throat (causing difficulty in swallowing or breathing)
- rapid heart rate, low blood pressure, collapse
- also, there may be skin reaction (including hives, rash and itching) – very rarely these can be severe with blistering and peeling skin.

Stomach and bowel problems including:

- indigestion or heartburn
- pains in your stomach (abdomen) or other abnormal stomach problems
- pass blood in your stools or motions
- pass black tarry stools or motions
- vomit any blood or dark particles that look like coffee grounds
- worsening of bowel problems (ulcerative colitis or Crohn's disease).

Blood disorders which can cause:

- unexplained or unusual bruising or bleeding
- sore throat or mouth ulcers
- fever (high temperature)
- extreme paleness or weakness and exhaustion.

Heart and circulation effects: Medicines such as Ibuprofen 200mg Tablets may be associated with a small increased risk of heart attack (myocardial infarction), high blood pressure, heart failure, swelling due to fluid build-up (oedema) or stroke.

Other side effects that may occur include:

Other stomach and intestine effects including: diarrhoea, flatulence, constipation and vomiting.

Liver problems that might be indicated by yellowing of the skin and eyes (jaundice) and/or pale coloured stools and dark urine.

Kidney problems that might be indicated by passing less or more urine than normal, cloudy urine, blood in the urine, pain in the back and/or swelling (particularly of the legs) – very rarely kidney failure.

Nervous system problems including headache and aseptic meningitis (which can have symptoms such as severe headache, stiff neck, disorientation, fever and eye sensitivity to light in those with existing auto-immune disorders such as lupus).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard.

5. STORING THE MEDICINE

Do not use after the expiry date shown on the pack.

Do not store above 25°C.

Store in the original container.

Keep out of the sight and reach of children.

6. FURTHER INFORMATION

What is in this medicine:

The active ingredient is: Ibuprofen 200mg per coated tablet.

The other ingredients are: Lactose, Maize Starch, Hypromellose, Sodium Starch Glycollate, Colloidal Anhydrous Silica, Magnesium Stearate, Sucrose, Talc, Titanium Dioxide (E171) and Carnauba Wax.

What this medicine looks like and contents of the pack

Ibuprofen 200mg Tablets are white, circular, sugar coated tablets with no markings.

This pack contains 6, 8, 12 or 16 tablets (not all pack sizes may be marketed).

Marketing Authorisation Holder:

Galpharm Healthcare Ltd., Wraffton, Braunton, Devon, EX33 2DL, United Kingdom.

Manufacturer:

Galpharm International Ltd., Elmhirst Park, Middle Field Road, Barnsley, South Yorkshire, S75 4LS, United Kingdom.

Text revised: December 2014.

Component code

Artwork reference: BTC206217