

FOR FAST RELIEF OF:

Headache	✓
Migraine	✓
Toothache	✓
Period pain	✓

FOR RELIEF OF THE SYMPTOMS OF:

Cold, flu, sore throat	✓
Rheumatic pain	✓
Lumbago	✓
Sciatica and neuralgia	✓



Directions: Dissolve tablets in water before taking.



Dosage: Adults 2-3 tablets every four hours.
Do not exceed 13 tablets in 24 hours.

If symptoms persist consult your doctor.

Do not exceed the stated dose.

DO NOT TAKE IF: you have, or have had, a stomach ulcer or haemophilia.

WARNINGS AND INTERACTIONS: Consult your doctor before taking if: you are asthmatic, allergic to aspirin, receiving regular medical treatment, such as medicines for blood clots or gout, or are pregnant or breastfeeding.

SIDE EFFECTS: In people sensitive to aspirin, reactions such as an asthma attack and skin rashes may occur occasionally.

Use may also induce gastrointestinal haemorrhage (vomiting of blood or passing tarry stools), occasionally major. Tell your doctor or pharmacist if you have any side effects after taking this product.

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN

Do not give to children aged under 16 years, unless on the advice of a doctor. Contains aspirin.

There is a possible association between aspirin and Reye's syndrome when given to children. Reye's syndrome is a very rare disease, which can be fatal. For this reason aspirin should not be given to children aged under 16 years, unless on the advice of a doctor.

Each tablet contains: Aspirin Ph. Eur.
300mg, calcium carbonate, maize starch, citric acid anhydrous, talc, sodium lauryl sulphate, saccharin, crospovidone and lime flavour.

Store below 25°C in a dry place.

