

Boots Wind Relief Tablets

Aluminium Hydroxide, Magnesium Hydroxide, Simeticone

GENTLE ACTION

- ✓ **Fast effective relief**
- ✓ **For the pain and discomfort of trapped wind, bloating, fullness after food and indigestion**

36

Read all of this carton for full instructions.

What this medicine is for

This medicine contains a wind reliever and antacids, which can be used to relieve bloating and the pain of trapped wind.

It can also be used to relieve the symptoms of indigestion, heartburn and excess acid.

Before you take this medicine

X Do not take:

- **If you are allergic** to any of the ingredients
- **If you have low levels of phosphate in your blood**
- **If you have an intolerance to some sugars**, unless your doctor tells you to (this medicine contains sorbitol and sucrose)
- **If are pregnant or breastfeeding**, unless your doctor tells you to

! Talk to your pharmacist or doctor:

- If you have kidney problems
- If you take other medicines – tetracycline antibiotics, chloroquine (to prevent malaria), quinolone antibiotics (e.g. ciprofloxacin), penicillamine (for rheumatoid arthritis), phenothiazines (e.g. chlorpromazine for mental problems)

✓ How to take this medicine

Check the foil is not broken before use. If it is, do not take that tablet.

Age	How many to take	How often to take
Adults and children of 12 years and over	One or two tablets	After meals, at bedtime, or when you need to.
Children of 5 to 11 years	One tablet	After meals, at bedtime, or when you need to.

Suck or chew the tablets.

Do not give to children under 5 years.

If symptoms do not go away within 5 days talk to your doctor.

If you take too many tablets: You may feel sick or be sick. Talk to a pharmacist or doctor if this happens.

Possible side effects

Most people will not have problems, but some may get some of these:

- Occasionally diarrhoea or constipation

Reporting of side effects

If you get any of the side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

How to store this medicine

Keep all medicines out of the sight and reach of children.

Use by the date on the end flap of the carton.

Active ingredients

Each tablet contains Dried Aluminium Hydroxide Gel (calculated as Aluminium Oxide) 170 mg, Magnesium Hydroxide 164 mg, Simeticone 25 mg.

Also contains: sucrose, sorbitol (E420), maize starch, povidone, microcrystalline cellulose, magnesium stearate, peppermint oil.

PL 00014/0287

Text prepared 05/18

Manufactured for the Marketing Authorisation holder The Boots Company PLC Nottingham NG2 3AA by Famar AVE Anthoussa Attiki 15349 Greece

If you need more advice ask your pharmacist.

WBAG321685