PATIENT INFORMATION LEAFLET regurgitation, heartburn and indigestion. This product contains methyl (E218) and GAVISCON DOUBLE ACTION MINT which may occur, for example, following propyl (E216) parahydroxybenzoates which Sodium alginate meals or during pregnancy and for may cause allergic reactions (possibly Sodium bicarbonate symptoms of excess stomach acidity delayed). Calcium carbonate Taking Other Medicines (hyperacidity). Do not take this product within two hours of Before taking Gaviscon Double Action Please read this leaflet carefully before you Mint taking other medicines by mouth as it can take this medicine. If you are not sure about Do not take this product if: interfere with the action of some other anything ask your pharmacist or doctor. · You know you are allergic to any of the medicines. This is especially important if What is Gaviscon Double Action Mint and ingredients as very rarely difficulty in you are taking antihistamines, antibiotics what is it used for? breathing and skin rashes have occurred (see (tetracyclines and quinolones such as Gaviscon Double Action Mint is a further information for a full list of norfloxacin), iron preparation, antifungals combination of two antacids (calcium ingredients). such as ketoconazole, digoxin and carbonate and sodium bicarbonate) and an Take special care before treatment with beta-blockers (for heart conditions). alginate and works in two ways: Gaviscon Double Action Mint: neuroleptics (for mental illness), thyroxine, 1) Neutralising excess stomach acid to This medicine contains small amounts of chloroquine (for malaria) and relieve the pain and discomfort. sodium (11.06mmol per 20ml) and calcium biphosphonates (for osteoporosis). (2) Forming a protective barrier over the (6.5 mmol per 20ml). If you have been How to take Gaviscon Double Action Mint stomach contents to soothe the burning pain Check that the cap seal is unbroken before advised to follow a diet restricted in either of in your chest. first using this product. these please consult your doctor. Gaviscon Double Action Mint is used for the You can take this product if you are For oral use. Shake well before use. treatment of symptoms of pregnant or breast feeding. gastro-oesophageal reflux such as acid 3020672 Adults and children 12 years and over: and consult your doctor immediately. Also (E218) and propyl (E216) Take 10-20ml (two to four 5ml spoonfuls) you can help to make sure that medicines parahydroxybenzoates, sodium saccharin, after meals and at bedtime, up to four times a remain as safe as possible by reporting any mint flavour, sodium hydroxide and water. unwanted side effects via the internet at day, or as directed by your doctor. Children This product does not contain sugar or under 12 years: Should only be taken on http://www.mhra.gov.uk/yellowcard gluten. medical advice. If you take too much of this Alternatively you can call Freephone Manufacturer and PL Holder in UK: product you may feel bloated. It is unlikely 0808 100 3352 (available from 10 a.m. to Reckitt Benckiser Healthcare (UK) Ltd., 2 p.m. Mondays to Fridays) or fill in a paper to cause you any harm, but please consult Hull, HU8 7DS. your doctor or pharmacist. If you forget a form available from your local pharmacy. Export distributors: Reckitt & Colman dose it is not necessary to double the dose How to store Gaviscon Double Action Mint (Overseas) Ltd., Hull, HU8 7DS, UK. next time, just carry on taking as before. Do not use this product after the expiry date Gaviscon and the sword and circle are trade After taking Gaviscon Double Action (EXP: month/year) shown marks. Mint: If symptoms persist after 7 days · Use within 6 months of opening Text Revised: July 2015 consult your doctor. · Keep out of the reach and sight of children Possible side effects Do not store above 30°C. Very rarely (less than 1 in 10,000 patients Do not refrigerate or freeze treated) an allergic reaction to the ingredients Further information Each 10 ml of oral suspension contains may occur. Symptoms of this may include skin rash, itching, difficulty breathing, 500mg sodium alginate, 213mg sodium bicarbonate and 325mg calcium carbonate as dizziness, or swelling of the face, lips,

the active ingredients.

any other side-effects stop taking the product. The other ingredients are carbomer, methyl

itongue or throat. If you experience these or