











Manufacturer and Product Usence Holber:
Reckits Benchiser Healthcare IIIC Limited, Hull, HUB 7DS.
Date of paramaters March 2013.
Gavincon Advance and the sword and circle are trademarks.

CUSTOMER INFO:
Minimum Point Size = 7.00pt
Leaflet Point Size = 9pt



Gaviscon Advance Aniseed flavour Oral Suspension. Sodium alginate and potassium hydrogen carbonate. Read all of this leaflet carefully before you start using this medicine because it contains important

- information for you.

  Keep this leaflet, You may need to read it again,
- Ask your pharmacist if you need more information or
- advice.

  If you get any side effects, talk to your pharmacist.

  This includes any possible side effects not listed in this leaflet. See section 4. You must talk to your doctor if you do not feel better or i you feel worse after 7 days

  In this baselet.

  1. What is this medicine and what is it used for?

  2. What you need to know before you take this medicine.

- 3. How to take this medicine.
- Possible side effects. 5. How to store this medicine
- 6. Contents of the pack and other information.

What is this medicine and what is if used for? Gaviscon Advance Aniseed flavour Oral Suspension belongs to a group of medicines caled reflux. suppressants. This product forms a protective layer that floats on top of the stomach contents. This layer prevents

floats on top of the stamech contents. This layer prevents relix and leeges for stomech contents save yfrom the lining of the food pipe to releve the symptoms of horathrum and seil midgestein. The can be be used to releve the symptoms of conditions such as histus hemia (protrusion of massle through an muscle walll, reflux coespitaglist inflamed food pipe) and symptoms of horasseness and other voice desorties, one throat and cough associated with reflux. It can also be taken to control hearthum symptoms which may occur when straing or following withdrawed of, medication treduce stomach and such as Proton Pump Inhibitors (PPTs) or FLA missporiss.

2. What you need to know before taking this medicine
Do not take Gaviscon Advance Aniseed flavour Oral

- If you know you are allergic (hypersensitive) to any of

the ingredients of in this product (Isted in section 6 for a full list of ingredients).

Warnings and precautions:
This medicine contains sodium (4,5 mmol per 10 mll, potassium (2,0 mmol per 10 ml) and calcium.

11 you have been advised to follow a diet restricted in any of these please consult your doctor before taking this

product.
Please also talk to your doctor regarding these salt
contents if you suffer or have suffered from significant
kidney or heart disease, as certain salts could interfere
with these diseases.
Taking other medicines:
Please tell your doctor or pharmacist if you are taking or
heart designed to the production of the plant of

Prease tell your octoor of paramacist if you are taxin have recently tacker or pharmacistines, including medicines obtained without prescription Pregnancy and breast-feeding: You can take this product if you are pregnant or breast-feeding. As with all medicines, the treatment duration should be limited as much as possible.

Important information about some of the ingredients of Genisson Advance Aniseed flavour Oral Suspension: This product contains methy (12:18) and propy (12:16) pare-hydroxyberusoites which may cause altergic reactions (possibly delayed). Driving and using machines: This medicine has no influence on the ability to drive and use machines.

3. How to take this medicine.

Deack that the can seal is unknoken before first using this

Check that the cap seal is unbroken before first using this product. Shake well before use, **Adults including the** product. Shake well before use, Anulus motioning the elderly and children 12 years and over. 5-10 ml (one to two 5ml spoonfuls) after meals and at bedime. Children under 12 years: Should only be taken on medical advice. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before. If you take too much of this product you may led bloaded it is unifiely to cause you any harm, but glosse consult your doctor or harmenist if this chosen't no away. pharmacist if this doesn't go away.

If symptoms persist after 7 days consult your doctor.

4. Possible side effects

Very rarely (less than 1 in 10,000 patients) there is a chance of an allergic reaction to the ingredients. Symptoms of this may include skin rash, taching, difficulty breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor mendiately. If you get any side effects face is, alto your doctor, pharmacist or nurse. This includes any possible side effects may lead in this leaflet, You can also report side effects directly via the fellow Card Scheme at http://www.unfra.gou.ulv/get/oucard. 5 y reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store this medicine.

safety of this medicine.

A. Hown to store this medicine
Do not use this product after the expiry date
(EXP: month) year) shown. Keep out of the sight and
reach of childre.
Do not rehignerate.
Medicines should not be disposed of via wastewater or
household waste. Ask your pharmacist how to dispose of

medicines no longer required. These measures will help o protect the environment.

to protect the environment. As Contents of the pack and other information. Each 10 ml of oral suspension contains 1000 mg sodium adjinate and 200 mg potassium hydrogen carbonate as the active imprediates are calcium carbonate, carbonner, methyl (E218) and propyl (E218) parahydroxybenozetis, sodium sacchain s, sodium sacchain s, sodium sacchain s, sodium sacchain s, sodium tente and purified valvate. This product does not contain suger or intent in this medicine is suitable in brates in suitable in the size in suitable in brates in suitable in the size in suitable in size in suitable in the size in s gluten, this medicine is available in bottles of 80ml, 100ml, 125ml, 140ml, 150ml, 180ml, 200ml, 250ml and

80m, 10mm, 12mm, 140mm, 180mm, 180mm, 200mm, 200mm.
Not all pack sizes may be marketed.
Manufacturer and Product Licence Holder:
Reckit Benobiser Healthcare (UK) Limited,
Hull, Hulb 70'S.
Date of March 2017.
Gaviscon Advance and the sword and circle are

trademarks.

**CUSTOMER INFO:** Minimum Point Size = 7.00pt Leaflet Point Size = 9pt



Gaviscon Advance Aniseed flavour Oral Suspension. Sodium alginate and potassium hydrogen carbonate. Read all of this leaflet carefully before you start using this medicine because it contains important

- information for you.

  Keep this leaflet, You may need to read it again,
- Ask your pharmacist if you need more information or
- advice.

  If you get any side effects, talk to your pharmacist.

  This includes any possible side effects not listed in this leaflet. See section 4. You must talk to your doctor if you do not feel better or i you feel worse after 7 days

  In this baselet.

  1. What is this medicine and what is it used for?

  2. What you need to know before you take this medicine.

- 3. How to take this medicine. Possible side effects.
- 5. How to store this medicine
- 6. Contents of the pack and other information.

 What is this medicine and what is if used for? Gaviscon Advance Aniseed flavour Oral Suspension belongs to a group of medicines caled reflux. suppressants. This product forms a protective layer that floats on top of the stomach contents. This layer prevents eflux and keeps the stomach contents away from the

reflux and keeps the stomach contents away from the inling of the Good piec to releve the symptoms of hearthurn and acid indigestion. It can also be used to hearthurn and acid indigestion. It can also be used to the symptoms of conditions such as histus hernia (protrusion of musuled through a musule wall, reflux esesphagits inflament food pip and symptoms of horsansess and other voice disorders, sore throat and cough associated with reflux, it can also be taken to control. hearthurn symptoms within may occur when taking, or following withdrawal of, medication to reduce stomach and such as Proton Pump Inhibitors (PPTs) or HZA masoniss.

2. What you need to know before taking this medicine
Do not take Gaviscon Advance Aniseed flavour Oral

- If you know you are allergic (hypersensitive) to any of

the ingredients of in this product (Issted in section 6 for a full list of ingredients).

Warnings and precautions:

Warnings and precautions:

This medicine contains sodium (4.6 mmol per 10 ml), potassium (2.0 mmol per 10 ml) and calcium.

- If you have been advised to follow a diet restricted in

ny of these please consult your doctor before taking this

product.
Please also talk to your doctor regarding these salt
contents if you suffer or have suffered from significant
kidney or heart disease, as certain salts could interfere
with these diseases.
Taking other medicines:
Please tell your doctor or pharmacist if you are taking or
heart designed to the production of the plant of

Prease tell your octoor of paramacist if you are taxin have recently tacker or pharmacistines, including medicines obtained without prescription Pregnancy and breast-feeding: You can take this product if you are pregnant or breast-feeding. As with all medicines, the treatment duration should be limited as much as possible.

Important information about some of the ingredients of Genisson Advance Aniseed flavour Oral Suspension: This product contains methy (12:18) and propy (12:16) pare-hydroxyberusoites which may cause altergic reactions (possibly delayed). Driving and using machines: This medicine has no influence on the ability to drive and use machines.

3. How to take this medicine.

Deack that the can seal is unknoken before first using this

Check that the cap seal is unbroken before first using this product. Shake well before use. Adults including the

product. Shake well before use, Anulus motioning the elderly and children 12 years and over. 5-10 ml (one to two 5ml spoonfuls) after meals and at bedime. Children under 12 years: Should only be taken on medical advice. If you forget a dose it is not necessary to double the dose next time, just carry on taking as before. If you take too much of this product you may led bloaded it is unifiely to cause you any harm, but glosse consult your doctor or harmenist if this chosen't no away. pharmacist if this doesn't go away.

If symptoms persist after 7 days consult your doctor.

4. Possible side effects

Very rarely (less than 1 in 10,000 patients) there is a chance of an alleging reaction to the ingredients. Symptoms of this may include shar rash, chining, difficulty breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor immediately. I you get any side effects hat to your doctor, pharmacist or nurse. This includes any possible side effects maked in this leafet, you can alway report side effects must listed in this leafet, you can alway report side effects must have provided to the side of this medicine.

5. How to store this medicine.

safety of this medicine.

A. Hown to store this medicine
Do not use this product after the expiry date
(EXP: month) year) shown. Keep out of the sight and
reach of childre.
Do not rehignerate.
Medicines should not be disposed of via wastewater or
household waste. Ask your pharmacist how to dispose of

medicines no longer required. These measures will help o protect the environment.

In protect the environment.

6. Contents of the pack and other information

6. Each 10 mf of oral suspension contains 100 mg sodium

alignate and 200 mg potassium hydrogen carbonate as

6. Each 10 mf of oral suspension contains 100 mg sodium

alignate and 200 mg potassium hydrogen carbonate as

6. Each 10 mf oral potassium hydrogen carbonate as

6. Each 10 mf oral potassium sachdarin, sodium

Apridrode, airsaed abour offerend from femal and

6. purified vaster. This product does not contain sugar or

6. and the time femilies is available in brother is medicine in sachdarin or available in brother. gluten, this medicine is available in bottles of 80ml, 100ml, 125ml, 140ml, 150ml, 180ml, 200ml, 250ml and

80m, 10mm, 12mm, 140mm, 180mm, 180mm, 200mm, 200mm.
Not all pack sizes may be marketed.
Manufacturer and Product Licence Holder:
Reckit Benobiser Healthcare (UK) Limited,
Hull, Hulb 70'S.
Date of March 2017.
Gaviscon Advance and the sword and circle are

trademarks.

TBR2138-D0006047 - English

GAVISCON ADVANCE ANISEED FLAVOUR ORAL SUSPENSION

**CUSTOMER INFO:** Minimum Point Size = 7.00pt Leaflet Point Size = 9pt















CUSTOMER INFO: Minimum Point Size = 7,00pt Leaflet Point Size = 9pt

T182117-01055166 - Englith

GAVISCON ADVANCE ANISEED FLAVOUR ORAL SUSPENSION