

Product Title: Tixylix Children's Cough Syrup	
Pack Size: 100ml	
Mat.No: 93040192112_100ML_PIL	
Dimensions: 210 x 148 flat (27 x 148 folded)	
Keyline Ref:	
Barcode:	
Pharmacode:	
Datamatrix: 3040192112	
SAP Code:	
Reason for Change: Change of manufacturer from BCM to T&R	
1. Colour: ■ Black	
2. Colour: ■ Keyline (Non-Printing)	
3. Colour: ■ Technical Info (Non-Printing)	
4. Colour: ■ Cirrus_Info_Box	
5. Colour:	
6. Colour:	
7. Colour:	
8. Colour:	
9. Colour:	
10. Colour:	
11. Colour:	
12. Colour:	
1. Fonts Used: Futura Std	
Point Size Used: Min:	Max: 9pt
2. Fonts Used:	
Point Size Used: Min:	Max:
3. Fonts Used:	
Point Size Used: Min:	Max:
4. Fonts Used:	
Point Size Used: Min:	Max:
5. Fonts Used:	
Point Size Used: Min:	Max:
Proof Number: 3	Date: 30.09.2022
Operator: EBR	Artwork Number: 1323148
CRN: MOC-2021-1195	Suffix: F
Local Authority submission required: Yes	
Manufacturer Art No: N/A	
Approved by:	Date:
APPROVED	
<i>By Katie Irving at 4:28 pm, Oct 17, 2022</i>	


Recto
14 mm

9 mm
14 mm

11.5 mm
14 mm


210 mm
14 mm

11.5 mm
4.2 mm



Children's Cough Syrup

Glycerol



PACKAGE LEAFLET: INFORMATION FOR THE USER
Read all of this leaflet carefully because it contains important information for you.
 This medicine is available without a prescription. However, you still need to give Tixylix Children's Cough Syrup carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your child's symptoms worsen or do not improve after 3 days.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Tixylix Children's Cough Syrup is and what it is used for
2. Before you give Tixylix Children's Cough Syrup
3. How to give Tixylix Children's Cough Syrup
4. Possible side effects
5. How to store Tixylix Children's Cough Syrup
6. Further information

1. WHAT TIXYLIX CHILDREN'S COUGH SYRUP IS AND WHAT IT IS USED FOR
 Tixylix Children's Cough Syrup is a soothing cough suppressant which relieves the symptoms of dry, tickly coughs. Each 5 ml dose of Tixylix Children's Cough Syrup contains 0.75 ml glycerol as the active ingredient. This medicine is intended for use in children aged 3 months to 5 years.

2. BEFORE YOU GIVE TIXYLIX CHILDREN'S COUGH SYRUP
Do not give Tixylix Children's Cough Syrup

- If your child is allergic (hypersensitive) to any of the ingredients of Tixylix Children's Cough Syrup (see Section 6).

Take special care with Tixylix Children's Cough Syrup

- Tixylix Children's Cough Syrup is not recommended for children under 3 months.

Taking other medicines
 Please tell your doctor or pharmacist if your child is taking, or has recently taken, any other medicines, including medicines obtained without a prescription. There are no known interactions between Tixylix Children's Cough Syrup and other medicines.

Taking Tixylix Children's Cough Syrup with food and drink
 Tixylix Children's Cough Syrup may be given with or without food.

Pregnancy and breast-feeding
 Ask your doctor or pharmacist for advice before taking any medicine. This medicine is intended for use in children. No information is available on the effect of glycerol taken during pregnancy or use during breast-feeding. Caution is therefore advised if taking Tixylix Children's Cough Syrup during pregnancy.

Driving and using machines
 This medicine is intended for use in children. Tixylix Children's Cough Syrup does not affect your ability to drive or operate machinery.

Important information about some of the ingredients of Tixylix Children's Cough Syrup.
 Tixylix Children's Cough Syrup contains:

- Maltitol, may have a mild laxative effect. Calorific value 2.3Kcal/g maltitol.
- Sodium benzoate (E211), 10mg per 5ml dose.
- Less than 1 mmol sodium (23mg) per 5ml dose, that is to say essentially "sodium-free"

3. HOW TO GIVE TIXYLIX CHILDREN'S COUGH SYRUP
 Always give Tixylix Children's Cough Syrup exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Verso
14 mm

9 mm
14 mm

11.5 mm
14 mm

210 mm
14 mm

11.5 mm
4.2 mm

Using the measuring spoon provided, the following doses may be given 3 to 4 times daily, by mouth, as required. Allow 3 to 4 hours between doses.

Age	Dose
Under 3 months	Do not give Tixylix Children's Cough Syrup
3 to 12 months	One 5 ml spoonful
1 to 5 years	Two 5 ml spoonfuls

After use, wipe the neck of the bottle clean and make sure the safety cap is properly closed. You can help by keeping your child warm and giving plenty of fluids. Make sure the bedroom is well ventilated but not cold. If your child's condition does not get better in 3 days, talk to your doctor.

If you give more Tixylix Children's Cough Syrup than you should
 Giving your child too much Tixylix Children's Cough Syrup may cause diarrhoea. If you have accidentally given your child more medicine than you should, or if you suspect that your child has taken extra medicine, please contact your doctor. Take along the bottle and the label, and any medicine that is left.

If you forget to give Tixylix Children's Cough Syrup
 Do not give your child a double dose to make up for a forgotten dose. If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS
 Like all medicines, Tixylix Children's Cough Syrup can cause side effects, although not everybody gets them. Tixylix Children's Cough Syrup may cause diarrhoea in overdose.

Reporting of side effects
 If your child gets any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk./yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE TIXYLIX CHILDREN'S COUGH SYRUP
 Keep out of the reach and sight of children. Do not store above 25°C. Keep the container tightly closed. Do not use Tixylix Children's Cough Syrup after the expiry date which is stated on the carton and the bottle label after <EXP>.

The expiry date refers to the last day of that month. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION
What Tixylix Children's Cough Syrup contains

- The active substance is glycerol (0.75 ml per 5ml dose).
- The other ingredients are liquid maltitol (lycasin), hydroxyethylcellulose, sodium benzoate (E211), citric acid monohydrate, sodium citrate, vanilla flavour (contains propylene glycol E1520), blackcurrant flavour (contains propylene glycol E1520), purified water.

What Tixylix Children's Cough Syrup looks like and contents of the pack
 Tixylix Children's Cough Syrup is a colour-free syrup. The following pack sizes are available:

- Amber-coloured glass bottle with tamper evident cap, containing 100ml. Measuring spoon provided.

Marketing Authorisation Holder and Manufacturer
 Thornton & Ross Ltd., Huddersfield, HD7 5QH, UK.
 For any information about this medicinal product, please contact the Marketing Authorisation Holder. This leaflet was last revised in October 2022

93040192112
