


Braille 25%



Varnish 25%



**What Peroxyl is used for:** Peroxyl is a medicated mouthwash used to relieve minor mouth and gum irritations by killing germs and/or preventing them from spreading and causing infection. When Peroxyl comes into contact with mouth tissue, oxygen is released in a bubbling action providing mechanical cleansing to flush out mouth debris and help in the relief of oral irritations.

**Before you use Peroxyl:** Do not use Peroxyl if you are allergic (hypersensitive) to hydrogen peroxide or any of the other ingredients of Peroxyl. You should speak to your dentist or doctor if you have a mouth irritation that persists beyond 7 days of treatment or is severe, if you have irritation due to the wearing of braces and/or dentures, if you develop swelling in the mouth or develop a fever.

**Special warnings:** Do not use in children under the age of 6 years. **DO NOT SWALLOW.** Avoid contact with eyes. Propylene glycol may cause skin irritation. **Pregnancy/Breast-Feeding:** Consult your doctor or dentist before use. Peroxyl contains levo-menthol, which if accidentally ingested in excessive amounts may cause fits (particularly in children or infants). If you, or anybody else using Peroxyl experience fits, seek medical help, always take this container with you.

**Directions for use:** Always use Peroxyl exactly as your dentist or pharmacist has told you. You should check with your dentist or pharmacist if you are not sure. You should not use Peroxyl at high doses or for a long time. The duration of the treatment should not exceed 7 days.

**Adults and children over 12 years:** Rinse with 10 ml (one capful, 10 ml line within cap) of Peroxyl for 1 minute, then spit out. Use 3 times daily, after meals and at bedtime, or as directed by your dentist or pharmacist.

**Children age 6-12 years:** As above but under the supervision of an adult.

**Possible side effects:** Like all medicines, Peroxyl can cause side effects, although not everybody gets them, these may include: minor irritation and swelling of the mouth, injury to the gums, black stain on the tongue and a feeling that your tongue is hairy. You are more likely to



# PEROXYL MOUTHWASH

with Hydrogen Peroxide



RELIEVES MINOR MOUTH AND GUM IRRITATIONS

Mint Flavour Antiseptic Action

experience these side effects if you use Peroxyl for a long time or at high doses. These side effects should disappear when you stop using Peroxyl. If you get any side effects, talk to your doctor, pharmacist or dentist. This includes any possible side effects not listed in this label. You can also report side effects directly via: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

**How to Store Peroxyl:** Keep out of the reach and sight of children, do not store above 25°C, do not use Peroxyl after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month. Do not use if you notice any signs of deterioration.

**What Peroxyl looks like and contains:** Peroxyl is a clear colorless liquid, available in 300 ml bottles. The active substance is hydrogen peroxide 1.5% w/v. The other ingredients are: Purified Water, Sorbitol 70%w/w (non-crystallising), Propylene glycol, Poloxamer 338, Polysorbate 20, Methyl salicylate, Levo-Menthol, Sodium saccharin.

**Marketing Authorisation Holder:** Colgate-Palmolive (UK) Ltd., Guildford, GU2 8JZ, UK.

**Manufacturer:** BCM Limited, Thane Road, Nottingham NG90 2PR, UK



Last revised in February 2021

5  
3  
8  
2  
10197585

300 ml e LOT:  
PL 00049/0038 EXP:

3  
4,5  
2

