

PACK OPS LYON GRAPHIC ARTWORK APPROVAL

<b>Code: 0 036 922905</b>	<b>VERSION</b> PTG : 0031 Version : F 180918	<b>COLORS</b>	
<b>PRODUCT:</b> WRA FTL PLUS CAT X3 RECTO	<b>INITIALS:</b> AJ - Composer ET/RH	<b>Noir/Black</b>	<b>Cold Foil</b>
<b>COUNTRY:</b> UK	<b>VERSION - DATE:</b> A 21/01/20 11h00 B 23/01/20 11h00 C 06/02/20 16h00 D 26/03/20 16h30	<b>P 2747</b>	
<b>Dimensions à plat (mm):</b> 147.2 x 16.6 x 111.6		<b>P 355</b>	
<b>Dimensions plié(e)s:</b> NA			
<b>PRINTED BASIS CODE (if applicable):</b> NA	<b>CANCEL AND REPLACE:</b> NA	Please use the official PANTONE® (P) matching system for an accurate color representation.	

**Contraindications:** In the absence of available data, the product should not be used on kittens less than 8 weeks old and/or weighing less than 1 kg. The product should not be used in ferrets less than 6 months old. Do not use on sick (e.g. systemic diseases, fever) or convalescent animals. **Do not use in rabbits, as adverse drug reactions and even mortality could occur.** In absence of studies, the use of the product is not recommended in non-target species. In the absence of safety studies, the minimum treatment interval is 4 weeks. Do not use in cases of hypersensitivity to the active substances or to any of the excipients. **Undesirable Effects:** Do not overdose. Among the very rare suspected adverse reactions, transient skin reactions at the application site (scaling, local hair loss, itching, redness) and general itching or hair loss have been reported after use. Excessive salivation, reversible nervous signs (increased sensitivity to stimulation, depression, other nervous signs) or vomiting have also been observed after use. If licking occurs, a brief period of excessive salivation may be observed due mainly to the nature of the carrier. The frequency of adverse reactions is defined using the following convention:  
- very common (more than 1 in 10 animals treated displaying adverse reaction(s))  
- common (more than 1 but less than 10 animals in 100 animals treated)  
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)  
- rare (more than 1 but less than 10 animals in 10,000 animals treated)  
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).  
If you notice any serious effects or other effects not mentioned here, please inform your veterinary surgeon. **Overdose:** The risk of experiencing adverse effects may however increase with overdosing (See Undesirable Effects). **Cats:** No undesirable effects were observed in target animal safety studies in cats and kittens aged 8 weeks and older and weighing about 1 kg treated once a month at five times the recommended dose for six consecutive months. Itching may occur following treatment. Overdose application of the product will cause a sticky appearance of hairs at the treatment spot. However, should this occur, it will disappear within 24 hours post application. **Ferrets:** In ferrets aged 6 months and older and treated once every 2 weeks for four treatments, at five times the recommended dose, body weight loss was observed in some animals.

**User warnings:** This product can cause mucous membrane, skin and eye irritation. Therefore, contact of the product with mouth, skin and eyes should be avoided. If the product is accidentally swallowed, seek medical advice immediately and show the package leaflet to the physician. Keep pipettes in original packaging until ready to use. In order to prevent children from getting access to used pipettes, dispose of used pipettes immediately. People with a known hypersensitivity (allergy) to insecticides or alcohol should avoid contact with the product. Avoid contents coming into contact with the fingers. If this occurs, wash hands with soap and water. After accidental exposure the eye should be rinsed carefully with plain water. Wash hands after use. Treated animals should not be handled until the application site is dry, and children should not be allowed to play with treated animals until the application site is dry. It is therefore recommended that animals are not treated during the day, but should be treated during the early evening, and that recently treated animals are not allowed to sleep with owners, especially children. Do not smoke, drink or eat during application. **Other precautions:** The alcohol carrier may have adverse effects on painted, varnished or other household surfaces or furnishings. **SPECIAL STORAGE CONDITIONS** Do not store above 30 °C. Store in the original package. Do not use this veterinary medicinal product after the expiry date stated on the package after EXP. The expiry date refers to the last day of the month. **SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS** Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements. Fipronil and (S)-methoprene may adversely affect aquatic organisms. Do not contaminate ponds, waterways or ditches with the product or empty containers. For animal treatment only. Keep out of the sight and reach of children. Blister card of 1 x 0.5 ml pipette  
Box of 1 blister card of 3 x 0.5 ml pipettes  
Box of 1 blister card of 4 x 0.5 ml pipettes  
Box of 2 blister cards of 3 x 0.5 ml pipettes  
Not all pack sizes may be marketed.

No print here except barcode+code as shown



922905

Batch:  
Exp.:

Do not store above 30 °C. Store in the original package. Do not use this veterinary medicinal product after the expiry date stated on the package after EXP.

The expiry date refers to the last day of the month. For animal treatment only. Keep out of the sight and reach of children.



# FRONTLINE

PLUS SPOT-ON CAT

Fipronil 10% w/v + (S)-methoprene 12% w/v SPOT-ON SOLUTION



For protection against fleas and ticks, and to prevent flea eggs from hatching, apply every 4 weeks.\*

### What does this product do?

- Kills fleas on your cat or ferret for 4 weeks.
- Prevents the development of flea eggs, larvae and pupae for 6 weeks when used in cats.
- Kills ticks on your cat for up to 2 weeks and on your ferret for up to 4 weeks.
- Kills biting lice on your cat.

### How to treat a home infestation:

In addition to treating your pet follow the 2 steps below:

- 1) Use a household flea treatment to kill eggs and larvae already in the home.
- 2) Regularly vacuum your carpets and furnishings.

For cats 8 weeks or older, weighing at least 1 kg and ferrets 6 months or older. Can be used on cats during pregnancy.

\* Tick efficacy for up to 2 weeks in cats. Minimum treatment interval 4 weeks. Before every use, read the information on the inside of the box for full instructions and all warnings.



**NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**  
Boehringer Ingelheim Animal Health UK Ltd, Bracknell, RG12 8YS, UK  
**MANUFACTURER**  
Boehringer Ingelheim Animal Health France SCS - 4 chemin du Calquet - 31000 Toulouse, France  
For customer support, email [petsUK@boehringer-ingelheim.com](mailto:petsUK@boehringer-ingelheim.com)

AVM-GSL  
Vm 08327/4267



3 FRONTLINE PLUS SPOT-ON CAT FOR CATS AND FERRETS

# FRONTLINE


PLUS SPOT-ON CAT

Fipronil 10% w/v + (S)-methoprene 12% w/v SPOT-ON SOLUTION



**KILLS FLEAS, TICKS & LICE**  
**PREVENTS THE DEVELOPMENT OF FLEA EGGS**

FOR **CATS AND FERRETS**



3 pipettes of 0.5 ml

**Marketing approval:**

COMPLETE NAME:  
DATE + SIGNATURE:

**Regulatory approval:**

COMPLETE NAME:  
DATE + SIGNATURE:

**NOT PRINTABLE / NON IMPRIMABLE**

**TECHNICAL NOTE(S) - INFO(S) TECHNIQUE**

- Unvarnished area / Réserve de vernis
- Artwork identification(s)
- Text areas / zones de textes
- Dieline / Découpe



6922905

# PACK OPS LYON GRAPHIC ARTWORK APPROVAL

<b>Code: 0 036 922905</b>	<b>VERSION</b> <span style="float: right;">PTG : 0031 Version : F 180918</span>	<b>COLORS</b>
<b>PRODUCT:</b> WRA FTL PLUS CAT X3 VERSO	<b>INITIALS:</b> AJ - Composer ET/RH <b>VERSION - DATE:</b> A 21/01/20 11h00 B 23/01/20 11h00 C 06/02/20 16h00 D 26/03/20 16h30	<b>Noir/Black</b>
<b>COUNTRY:</b> UK <b>Dimensions à plat (mm):</b> 147.2 x 16.6 x 111.6 <b>Dimensions plié(e)s:</b> NA	<b>CANCEL AND REPLACE:</b> NA	! Please use the official PANTONE® (P) matching system for an accurate color representation.
<b>PRINTED BASIS CODE (if applicable):</b> NA		

# FRONTLINE

## PLUS SPOT-ON CAT

Fipronil 10% w/v + (S)-methoprene 12% w/v

### STATEMENT OF ACTIVE AND OTHER SUBSTANCES

Clear amber solution containing:

Fipronil	50.00 mg
(S)-methoprene	60.00 mg
Butylhydroxyanisole (E320)	0.10 mg
Butylhydroxytoluene (E321)	0.05 mg
Excipient(s): q.s	0.5 ml

**PHARMACEUTICAL FORM:** Spot-on solution.

### INDICATION(S)

**In cats:**

To be used against infestations with fleas, alone or in association with ticks and/or biting lice.

- Treatment of flea infestations (*Ctenocephalides spp.*). Insecticidal efficacy against new infestations with adult fleas persists for 4 weeks. Prevention of the multiplication of fleas by inhibiting the development of eggs (ovicidal activity), larvae and pupae (larvicidal activity) originating from eggs laid by adult fleas for six weeks after application.

- Treatment of tick infestations (*Ixodes ricinus*, *Dermacentor variabilis*, *Rhipicephalus sanguineus*). The product has a persistent acaricidal efficacy for up to 2 weeks against ticks (based on experimental data).

- Treatment of infestations with biting lice (*Felicola subrostratus*).

**In ferrets:**

To be used against infestations with fleas, alone or in association with ticks.

- Treatment of flea infestations (*Ctenocephalides spp.*). Insecticidal efficacy against new infestations with adult fleas persists for 4 weeks. Prevention of the multiplication of fleas by inhibiting the development of eggs (ovicidal activity), larvae and pupae (larvicidal activity) originating from eggs laid by adult fleas.

- Treatment of tick infestations (*Ixodes ricinus*). The product has a persistent acaricidal efficacy for 4 weeks against ticks (based on experimental data).

Fipronil kills fleas within 24 hours and ticks and lice within 48 hours post-exposure.

### METHOD AND ROUTE(S) OF ADMINISTRATION

Spot-on

### METHOD OF ADMINISTRATION

1. Take the pipette out of its packaging.

Hold the pipette upright. Tap the narrow part of the pipette to ensure the contents remain within the main body of the pipette. Snap back the tip.

2. Part the coat on the back of the animal at the base of the neck in front of the shoulder blades until the skin is visible.

3. Place the tip of the pipette on the skin and squeeze the pipette several times to empty its contents completely and directly onto the skin in one spot.



### Dosage

One pipette of 0.5 ml per cat, corresponding to a minimum recommended dose of 5 mg/kg for fipronil and 6 mg/kg for (S)-methoprene, by topical application to the skin.

One pipette of 0.5 mL per ferret corresponding to a dose of 50 mg for fipronil and 60 mg for (S)-methoprene per ferret, by topical application to the skin.

### SPECIAL WARNING

For external use only.

### Use during pregnancy and lactation

The product can be used during pregnancy in cats. Consult your veterinary surgeon before using the product in lactating queens, or in pregnant or lactating jills.

### Precautions

Animals should be weighed accurately prior to treatment.

It is important to make sure that the product is applied to an area where the animal cannot lick it off and to make sure that animals do not lick each other following treatment.

When treating infestations of parasites, all in-contact animals should be treated with an appropriate product at the same time.

Do not use simultaneously with other flea products which are applied directly on to the animal.

No data on the effect of bathing/shampooing on the efficacy of the product in cats and ferrets are available. However, based on information available for dogs shampooed from 2 days after application of the product, it is not recommended to bath animals within 2 days after application of the product.

Avoid the contact with the animal's eyes.

There may be an attachment of single ticks. For this reason a transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.

Fleas from pets often infest the animal's basket, bedding and regular resting areas such as carpets and soft furnishings which should be treated, in case of massive infestation and at the beginning of the control measures, with a suitable insecticide and vacuumed regularly. The potential toxicity of the product for kittens of less than 8 weeks of age in contact with a treated queen is not documented. Special care should be taken in this case.

## Marketing approval:

COMPLETE NAME:

DATE + SIGNATURE:

## Regulatory approval:

COMPLETE NAME:

DATE + SIGNATURE:

NOT PRINTABLE  
NON IMPRIMABLE

### TECHNICAL NOTE(S) - INFO(S) TECHNIQUE

Unvarnished area / Réserve de vernis

Artwork identification(s)

Text areas / zones de textes

Dieline / Découpe

