

BRaille TEXT:

HIBISCRUB

Font: Marburg Medium 28,5 pt.
UK Braille



70mm

45mm

amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

What Hibiscrub looks like and contents of the pack
Hibiscrub is a clear, red, slightly viscous solution and comes in white bottles containing 125 ml, 250 ml, 500 ml or 5 litres of liquid.

Marketing Authorisation Holder
Regent Medical Limited, Medlock Street, Oldham, Lancashire, OL1 3HS, UK.

Manufacturer
BCM Ltd, Thane Road, Beeston, Nottingham, NG90 2PR, UK.

This leaflet was last revised in 06/2021.

Height - 70mm

Base Label
45mm

5. How to store Hibiscrub
Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is printed on the bottle after "EXP". The expiry date refers to the last day of that month. Do not store above 25°C. Store in the original package. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Hibiscrub contains

- The active substance is chlorhexidine gluconate 4% w/v (40 mg/ml)
- The other ingredients are poloxamer 237, isopropyl alcohol, lauryl dimethyl

HIBISCURB®
4% w/v Cutaneous Solution
Chlorhexidine gluconate 40 mg/ml
Antimicrobial Skin Cleanser for antiseptic handwashing on the ward, pre-operative surgical hand disinfection and patient pre- and post-operative skin antiseptics.

Read the package leaflet before use, which is given when the label is peeled back.

Excipients: poloxamer 237, isopropyl alcohol, lauryl dimethyl amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

Precautions: Avoid contact with eyes, brain, meninges, eyes and middle ear. Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns.

Keep out of the sight and reach of children. Do not store above 25°C. Store in the original package.

Hibiscrub® is a trade mark of Regent Medical Ltd.

7 332551 437299

PVEP4

FEEL HERE BUT DO NOT REMOVE

Height - 70mm

Page 16 - 34mm Front Cover - 42mm

Package leaflet: Information for the patient

HIBISCURB®
4% w/v Cutaneous Solution
Chlorhexidine gluconate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

(hives), skin irritation and blisters.

Inflammation of the membranes of the brain and spinal cord has been reported when they have come into direct contact with chlorhexidine.

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

Height - 70mm

Inside Cover Page 15

Hibiscrub is also used to disinfect hands after handling infected materials (antiseptic handwashing).

2. What you need to know before you use Hibiscrub

Do not use Hibiscrub

- if you are allergic (hypersensitive) to chlorhexidine gluconate or any of the other ingredients of this medicine (listed in section 6).
- Hibiscrub contains soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

Warnings and precautions
Talk to your doctor, pharmacist or nurse before using Hibiscrub.

- Avoid contact with the eyes, brain, meninges (the membranes surrounding the brain and spinal cord) and middle ear. If Hibiscrub comes into contact with the

If you accidentally swallow any Hibiscrub
You should go to your nearest Accident and Emergency department or contact your doctor immediately. Take this leaflet and any other packaging with you so they know what you have taken.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may occur; it is not known how often they occur but available data suggests these are very rare (probably affect fewer than 1 in every 10,000 people).

Height - 70mm

Page 4 - 34mm Page 13 - 34mm

- Serious allergic reactions which can cause breathing difficulties, weakness, collapse and death, and can sometimes be preceded by swelling (in particular lip/tongue and facial swelling), or rash.

Allergic reactions to chlorhexidine-impregnated patches have been reported rarely when used in newborn babies.

At the first sign of any of these reactions, tell your doctor, pharmacist or nurse and application of Hibiscrub should be stopped immediately.

Other possible side effects, for which it is not known how often they occur, are: allergic skin disorders such as dermatitis (inflammation of the skin), pruritus (itch), erythema (redness of the skin), eczema, rash, urticaria

What is in this leaflet

- What Hibiscrub is and what it is used for
- What you need to know before you use Hibiscrub
- How to use Hibiscrub
- Possible side effects
- How to store Hibiscrub
- Contents of the pack and other information

1. What Hibiscrub is and what it is used for

Hibiscrub belongs to a group of medicines called antiseptics. This means that it helps to prevent infections by killing germs on the skin.

Hibiscrub cutaneous solution contains the active ingredient chlorhexidine gluconate 4% w/v (40 mg/ml) which has antimicrobial properties. Hibiscrub is used to disinfect the hands and skin prior to or after an operation or clean procedure (pre- and post-operative hand and skin disinfection / antiseptics).

Height - 70mm

Page 14 Page 3

- Use 25 ml of Hibiscrub as a soap solution.
- Wash the whole body. Start with the face. Work down the body. Pay special attention to the skin around the nose, under the arms, around the belly button, between the legs and around the bottom.
- Rinse the whole body.
- Repeat the wash. Use the same amount of Hibiscrub. This time, wash the hair as well.
- Rinse the whole body and dry thoroughly with a clean towel.

For Preventing Infection After Your Operation
Wash the whole of your body, excluding the operation site, in the bath or the shower. This is usually done on the third day after the operation. Use the method described above.

eyes, wash out immediately and thoroughly with water.

- In patients with head or spinal injuries or a perforated ear drum (an ear drum with a hole or tear in it), the benefit of use in pre-operative preparation should be evaluated against the risk of contact.
- Common bleaches (which contain hypochlorites) may cause brown stains to develop on fabrics which have previously been in contact with chlorhexidine.
- Hibiscrub's action may be reduced if used with soap or detergents.
- Hibiscrub is flammable. Do not use with ignition sources until dry.
- If you go into hospital, let your doctor know if you are using Hibiscrub.

Children
Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns.

Height - 70mm

Page 12 - 34mm Page 5 - 34mm

Hibiscrub contains isopropyl alcohol and citrus perfume

This medicine contains 40 mg isopropyl alcohol in each ml. Dryness, irritation, redness or burning sensation of the skin may occur. Allergic contact dermatitis or contact allergies are very rare.

This medicine contains fragrance with 3-Methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one, Amyl cinnamal, Amylcinnamyl alcohol, Anisyl alcohol, Benzyl alcohol, Benzyl benzoate, Benzyl cinnamate, Benzyl salicylate, Cinnamal, Citral, Citronellol, Coumarin, d-Limonene, Eugenol, Farnesol, Geraniol, Hexyl cinnamaldehyde, Hydroxymethylpentyl-cyclohexenecarboxaldehyde, Isoeugenol, Lilial, Linalool, Methyl heptane carbonate, Oak moss and Tree moss. These may cause allergic reactions. The fragrance also contains

Continue washing hands and wrists until one minute has elapsed. Clean the fingernails with a brush or scraper. Rinse thoroughly. For pre-operative use repeat the wash a second time for two minutes.

The Hibiscrub handwashing technique is based on a procedure described by Ayliffe GAJ et al. J Clin Path 1978; 31: 923.

For Antiseptic Handwashing on the Ward
Wet the hands and forearms, apply 5 ml of Hibiscrub and wash for one minute. Rinse thoroughly and dry.

For Preventing Infection Before Your Operation
Wash the whole of your body in the bath or the shower on at least two occasions. This is usually done on the day before the operation and the day of the operation. Use the following method:

Height - 70mm

Page 6 Page 11

cause local skin reactions (such as stinging or burning sensation) or increase skin reactions caused by other products when applied on the same area.

3. How to use Hibiscrub

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you. Check with your doctor, pharmacist or nurse if you are not sure.

FOR EXTERNAL USE ONLY

Do not inject. Do not use in the body's internal spaces.

Chlorhexidine must not be used in direct contact with nerve tissue, for example brain and spinal cord tissue.

For Pre-operative Surgical Hand Disinfection
Wet the hands and forearms.

Wash with 5 ml of Hibiscrub using the following procedure, each step consisting of five strokes backwards and forwards.

- Palm to palm.
- Right palm over left dorsum and left palm over right dorsum.
- Palm to palm fingers interlaced.

Height - 70mm

Page 8 - 34mm Page 9 - 34mm

- Back of fingers to opposing palms with fingers interlocked.
- Rotational rubbing of right thumb clasping in left palm and vice versa.
- Rotational rubbing backwards and forwards of tips of fingers and thumb of right hand in left palm and vice versa.

Bergamot oil (containing bergapten), Soya oil, Ethanol, Methyl benzoate, Benzoic acid, Balsam of Peru, Formaldehyde and Sodium laurilsulfate.

Bergamot oil (containing bergapten) may increase sensitivity to UV light (natural and artificial sunlight).

Ethanol may cause burning sensation on damaged skin.

Benzoic acid, Benzyl benzoate and Methyl benzoate may cause local irritation. Benzoic acid, Benzyl benzoate and Methyl benzoate may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old).

Balsam of Peru may cause skin reactions.

Formaldehyde may cause local skin reactions (e.g. contact dermatitis).

Sodium laurilsulfate may

Height - 70mm

Page 10 Page 7


BRAILLE TEXT:

HIBISCRUB

Font: Marburg Medium 28,5 pt.
EU Braille

H I B I S C R U B

58mm



85mm

Height - 85mm

Marketing Authorisation Holder
Regent Medical Limited, Medlock Street, Oldham, Lancashire, OL1 3HS, UK.

Manufacturer
BCM Ltd., Thane Road, Beeston, Nottingham, NG90 2PR, UK.

This leaflet was last revised in 06/2021.

Base Label
58mm

Height - 85mm

6. Contents of the pack and other information

What Hibiscrub contains

- The active substance is chlorhexidine gluconate 4% w/v (40 mg/ml)
- The other ingredients are poloxamer 237, isopropyl alcohol, lauryl dimethyl amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

What Hibiscrub looks like and contents of the pack

Hibiscrub is a clear, red, slightly viscous solution and comes in white bottles containing 125 ml, 250 ml, 500 ml or 5 litres of liquid.

HIBISCRUB®
4% w/v Cutaneous Solution
Chlorhexidine gluconate 40 mg/ml

Antimicrobial Skin Cleanser for antiseptic handwashing on the ward, pre-operative surgical hand disinfection and patient pre- and post-operative skin antiseptics.

Read the package leaflet before use, which is given when the label is peeled back.

Excipients: poloxamer 237, isopropyl alcohol, lauryl dimethyl amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

Precautions: Avoid contact with eyes, brain, meninges, eyes and middle ear. Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns.

FOR EXTERNAL USE ONLY
Keep out of the sight and reach of children. Do not store above 25°C. Store in the original package.
Regent Medical Limited, Medlock Street, Oldham, Lancashire, OL1 3HS, UK.

PL 22099/0001
Hibiscrub® is a trade mark of Regent Medical Ltd.

5 060097 930784 >

PEEL HERE BUT DO NOT REMOVE

Page 16 - 47mm Front Cover - 55mm

Package leaflet: Information for the patient

HIBISCRUB®
4% w/v Cutaneous Solution
Chlorhexidine gluconate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Hibiscrub

Keep this medicine out of the sight and reach of children.
Do not use this medicine after the expiry date which is printed on the bottle after "EXP". The expiry date refers to the last day of that month.
Do not store above 25°C. Store in the original package.
Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Inside Cover Page 15

Height - 85mm

and post-operative hand and skin disinfection / antiseptics).

Hibiscrub is also used to disinfect hands after handling infected materials (antiseptic handwashing).

2. What you need to know before you use Hibiscrub

Do not use Hibiscrub

- if you are allergic (hypersensitive) to chlorhexidine gluconate or any of the other ingredients of this medicine (listed in section 6).
- Hibiscrub contains soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using Hibiscrub.

- Avoid contact with the eyes, brain, meninges (the membranes surrounding the brain and spinal cord) and middle ear. If Hibiscrub

If you accidentally swallow any Hibiscrub

You should go to your nearest Accident and Emergency department or contact your doctor immediately. Take this leaflet and any other packaging with you so they know what you have taken.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may occur; it is not known how often they occur but available data suggests these are very rare (probably affect fewer than 1 in every 10,000 people).

- Serious allergic reactions which can cause breathing difficulties, weakness, collapse and death, and

Page 4 - 47mm Page 13 - 47mm

can sometimes be preceded by swelling (in particular lip/tongue and facial swelling), or rash.

Allergic reactions to chlorhexidine-impregnated patches have been reported rarely when used in newborn babies.

At the first sign of any of these reactions, tell your doctor, pharmacist or nurse and application of Hibiscrub should be stopped immediately.

Other possible side effects, for which it is not known how often they occur, are: allergic skin disorders such as dermatitis (inflammation of the skin), pruritus (itch), erythema (redness of the skin), eczema, rash, urticaria (hives), skin irritation and blisters. Inflammation of the membranes of the brain and spinal cord has been reported when they have come into direct contact with chlorhexidine.

What is in this leaflet

- What Hibiscrub is and what it is used for
- What you need to know before you use Hibiscrub
- How to use Hibiscrub
- Possible side effects
- How to store Hibiscrub
- Contents of the pack and other information

1. What Hibiscrub is and what it is used for

Hibiscrub belongs to a group of medicines called antiseptics. This means that it helps to prevent infections by killing germs on the skin.

Hibiscrub cutaneous solution contains the active ingredient chlorhexidine gluconate 4% w/v (40 mg/ml) which has antimicrobial properties. Hibiscrub is used to disinfect the hands and skin prior to or after an operation or clean procedure (pre-

Page 14 Page 3

Height - 85mm

- Use 25 ml of Hibiscrub as a soap solution.
- Wash the whole body. Start with the face. Work down the body. Pay special attention to the skin around the nose, under the arms, around the belly button, between the legs and around the bottom.
- Rinse the whole body.
- Repeat the wash. Use the same amount of Hibiscrub. This time, wash the hair as well.
- Rinse the whole body and dry thoroughly with a clean towel.

For Preventing Infection After Your Operation

Wash the whole of your body, excluding the operation site, in the bath or the shower. This is usually done on the third day after the operation. Use the method described above.

comes into contact with the eyes, wash out immediately and thoroughly with water.

- In patients with head or spinal injuries or a perforated ear drum (an ear drum with a hole or tear in it), the benefit of use in pre-operative preparation should be evaluated against the risk of contact.
- Common bleaches (which contain hypochlorites) may cause brown stains to develop on fabrics which have previously been in contact with chlorhexidine.
- Hibiscrub's action may be reduced if used with soap or detergents.
- Hibiscrub is flammable. Do not use with ignition sources until dry.
- If you go into hospital, let your doctor know if you are using Hibiscrub.

Children

Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns.

Page 12 - 47mm Page 5 - 47mm

Hibiscrub contains isopropyl alcohol and citrus perfume

This medicine contains 40 mg isopropyl alcohol in each ml. Dryness, irritation, redness or burning sensation of the skin may occur. Allergic contact dermatitis or contact allergies are very rare.

This medicine contains fragrance with 3-Methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one, Amyl cinnamal, Amylcinnamyl alcohol, Anisyl alcohol, Benzyl alcohol, Benzyl benzoate, Benzyl cinnamate, Benzyl salicylate, Cinnamal, Citral, Citronellol, Coumarin, d-Limonene, Eugenol, Farnesol, Geraniol, Hexyl cinnamaldehyde, Hydroxymethylpentyl-cyclohexenecarboxaldehyde, Isoeugenol, Linal, Linalool, Methyl heptane carbonate, Oak moss and Tree moss. These may cause allergic reactions. The fragrance also contains Bergamot oil (containing bergapten), Soya oil,

Continue washing hands and wrists until one minute has elapsed. Clean the fingernails with a brush or scraper. Rinse thoroughly. For pre-operative use repeat the wash a second time for two minutes.

The Hibiscrub handwashing technique is based on a procedure described by Ayliffe GAJ et al. J Clin Path 1978; 31: 923.

For Antiseptic Handwashing on the Ward

Wet the hands and forearms, apply 5 ml of Hibiscrub and wash for one minute. Rinse thoroughly and dry.

For Preventing Infection Before Your Operation

Wash the whole of your body in the bath or the shower on at least two occasions. This is usually done on the day before the operation and the day of the operation. Use the following method:

Page 6 Page 11

Height - 85mm

reactions caused by other products when applied on the same area.

3. How to use Hibiscrub

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you. Check with your doctor, pharmacist or nurse if you are not sure.


FOR EXTERNAL USE ONLY

Do not inject. Do not use in the body's internal spaces.


Chlorhexidine must not be used in direct contact with nerve tissue, for example brain and spinal cord tissue.

For Pre-operative Surgical Hand Disinfection


Wet the hands and forearms. Wash with 5 ml of Hibiscrub using the following procedure, each step consisting of five strokes backwards and forwards.



1. Palm to palm.




2. Right palm over left dorsum and left palm over right dorsum.




3. Palm to palm fingers interlaced.


Page 8 - 47mm Page 9 - 47mm



4. Back of fingers to opposing palms with fingers interlocked.



5. Rotational rubbing of right thumb clasping in left palm and vice versa.



6. Rotational rubbing backwards and forwards of tips of fingers and thumb of right hand in left palm and vice versa.

Ethanol, Methyl benzoate, Benzoic acid, Balsam of Peru, Formaldehyde and Sodium laurilsulfate.

Bergamot oil (containing bergapten) may increase sensitivity to UV light (natural and artificial sunlight).

Ethanol may cause burning sensation on damaged skin.

Benzoic acid, Benzyl benzoate and Methyl benzoate may cause local irritation. Benzoic acid, Benzyl benzoate and Methyl benzoate may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old).

Balsam of Peru may cause skin reactions.

Formaldehyde may cause local skin reactions (e.g. contact dermatitis). Sodium laurilsulfate may cause local skin reactions (such as stinging or burning sensation) or increase skin

Page 10 Page 7


BRAILLE TEXT:

HIBISCRUB










Font: Marburg Medium 28,5 pt.
UK Braille



45mm



115mm



BRaille TEXT:

HIBISCRUB

Font: Marburg Medium 28,5 pt.
UK Braille

H I B I S C R U B

HIBISCRUB®

4% w/v Cutaneous Solution
Chlorhexidine gluconate 40 mg/ml

H I B I S C R U B



150mm

WZBA7

5l e



102mm

Height - 150mm

5. How to store Hibiscrub

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is printed on the bottle after "EXP". The expiry date refers to the last day of that month. Do not store above 25°C. Store in the original package. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Hibiscrub contains

- The active substance is chlorhexidine gluconate 4% w/v (40 mg/ml)
- The other ingredients are poloxamer 237, isopropyl alcohol, lauryl dimethyl amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

What Hibiscrub looks like and contents of the pack

Hibiscrub is a clear, red, slightly viscous solution and comes in white bottles containing 125 ml, 250 ml, 500 ml or 5 litres of liquid.

Marketing Authorisation Holder

Regent Medical Limited, Medlock Street, Oldham, Lancashire, OL1 3HS, UK.

Manufacturer

BCM Ltd., Thane Road, Beeston, Nottingham, NG90 2PR, UK.

This leaflet was last revised in 06/2021.

Base Label - 102mm

Package leaflet: Information for the patient

HIBISCRUB®

4% w/v Cutaneous Solution
Chlorhexidine gluconate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you.
- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Hibiscrub is and what it is used for
2. What you need to know before you use Hibiscrub
3. How to use Hibiscrub
4. Possible side effects
5. How to store Hibiscrub
6. Contents of the pack and other information

1. What Hibiscrub is and what it is used for

Hibiscrub belongs to a group of medicines called antiseptics. This means that it helps to prevent infections by killing germs on the skin.

Hibiscrub cutaneous solution contains the active ingredient chlorhexidine gluconate 4% w/v (40 mg/ml) which has antimicrobial properties.

Hibiscrub is used to disinfect the hands and skin prior to or after an operation or clean procedure (pre- and post-operative hand and skin disinfection / antiseptis).

Hibiscrub is also used to disinfect hands after handling infected materials (antiseptic handwashing).

2. What you need to know before you use Hibiscrub

Do not use Hibiscrub

- if you are allergic (hypersensitive) to chlorhexidine gluconate or any of the other ingredients of this medicine (listed in section 6).
- Hibiscrub contains soya oil. If you are allergic to peanut or soya, do not use this medicinal product.

Warnings and precautions

- Talk to your doctor, pharmacist or nurse before using Hibiscrub.
- Avoid contact with the eyes, brain, meninges (the membranes surrounding the brain and spinal cord) and middle ear. If Hibiscrub comes into contact with the eyes, wash out immediately and thoroughly with water.
- In patients with head or spinal injuries or a perforated ear drum (an ear drum with a hole or tear in it), the benefit of use in pre-operative preparation should be evaluated against the risk of contact.
- Common bleaches (which contain hypochlorites) may cause brown stains to develop on fabrics which have previously been in contact with chlorhexidine.
- Hibiscrub's action may be reduced if used with soap or detergents.
- Hibiscrub is flammable. Do not use with ignition sources until dry.
- If you go into hospital, let your doctor know if you are using Hibiscrub.

Children

Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns.

Hibiscrub contains isopropyl alcohol and citrus perfume

This medicine contains 40 mg isopropyl alcohol in each ml. Dryness, irritation, redness or burning sensation of the skin may occur. Allergic contact dermatitis or contact allergies are very rare.

This medicine contains fragrance with

3-Methyl-4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-one, Amyl cinnamal, Amylcinnamyl alcohol, Anisyl alcohol, Benzyl alcohol, Benzyl benzoate, Benzyl cinnamate, Benzyl salicylate, Cinnamal, Citral, Citronellol, Coumarin, d-Limonene, Eugenol, Farnesol, Geraniol, Hexyl cinnamaldehyde, Hydroxymethylpentyl-cyclohexenecarboxaldehyde, Isoeugenol, Linalol, Linalool, Methyl heptine carbonate, Oak moss and Tree moss. These may cause allergic reactions. The fragrance also contains Bergamot oil (containing bergapten), Soya

oil, Ethanol, Methyl benzoate, Benzoic acid, Balsam of Peru, Formaldehyde and Sodium laurilsulfate.

Bergamot oil (containing bergapten) may increase sensitivity to UV light (natural and artificial sunlight).

Ethanol may cause burning sensation on damaged skin.

Benzoic acid, Benzyl benzoate and Methyl benzoate may cause local irritation. Benzoic acid, Benzyl benzoate and Methyl benzoate may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old).

Balsam of Peru may cause skin reactions. Formaldehyde may cause local skin reactions (e.g. contact dermatitis).

Sodium laurilsulfate may cause local skin reactions (such as stinging or burning sensation) or increase skin reactions caused by other products when applied on the same area.

3. How to use Hibiscrub

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you. Check with your doctor, pharmacist or nurse if you are not sure.

FOR EXTERNAL USE ONLY

Do not inject. Do not use in the body's internal spaces.

Chlorhexidine must not be used in direct contact with nerve tissue, for example brain and spinal cord tissue.

For Pre-operative Surgical Hand Disinfection

Wet the hands and forearms. Wash with 5 ml of Hibiscrub using the following procedure, each step consisting of five strokes backwards and forwards.


Height - 150 mm

Inside Cover


Page 3

Page 4


Height - 150 mm




1. Palm to palm.




2. Right palm over left dorsum and left palm over right dorsum.




3. Palm to palm fingers interlaced.



4. Back of fingers to opposing palms with fingers interlocked.



5. Rotational rubbing of right thumb clasping in left palm and vice versa.



6. Rotational rubbing backwards and forwards of tips of fingers and thumb of right hand in left palm and vice versa.

Continue washing hands and wrists until one minute has elapsed. Clean the fingernails with a brush or scraper. Rinse thoroughly. For pre-operative use repeat the wash a second time for two minutes.

The Hibiscrub handwashing technique is based on a procedure described by Ayliffe GAJ et al. J Clin Path 1978; 31: 923.

For Antiseptic Handwashing on the Ward

Wet the hands and forearms, apply 5 ml of Hibiscrub and wash for one minute. Rinse thoroughly and dry.

For Preventing Infection Before Your Operation

Wash the whole of your body in the bath or the shower on at least two occasions. This is usually done on the day before the operation and the day of the operation. Use the following method:

- Use 25 ml of Hibiscrub as a soap solution.
- Wash the whole body. Start with the face. Work down the body. Pay special attention to the skin around the nose, under the arms, around the belly button, between the legs and around the bottom.
- Rinse the whole body.
- Repeat the wash. Use the same amount of Hibiscrub. This time, wash the hair as well.
- Rinse the whole body and dry thoroughly with a clean towel.

For Preventing Infection After Your Operation

Wash the whole of your body, excluding the operation site, in the bath or the shower. This is usually done on the third day after the operation. Use the method described above.

If you accidentally swallow any Hibiscrub

You should go to your nearest Accident and Emergency department or contact your doctor immediately. Take this leaflet and any other packaging with you so they know what you have taken.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may occur; it is not known how often they occur but available data suggests these are very rare (probably affect fewer than 1 in every 10,000 people).

- Serious allergic reactions which can cause breathing difficulties, weakness, collapse and death, and can sometimes be preceded by swelling (in particular lip/tongue and facial swelling), or rash.

Allergic reactions to chlorhexidine-impregnated patches have been reported rarely when used in newborn babies.

At the first sign of any of these reactions, tell your doctor, pharmacist or nurse and application of Hibiscrub should be stopped immediately.

Other possible side effects, for which it is not known how often they occur, are: allergic skin disorders such as dermatitis (inflammation of the skin), pruritus (itch), erythema (redness of the skin), eczema, rash, urticaria (hives), skin irritation and blisters.

Inflammation of the membranes of the brain and spinal cord has been reported when they have come into direct contact with chlorhexidine.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

HIBISCRUB®

4% w/v Cutaneous Solution
Chlorhexidine gluconate 40 mg/ml

Antimicrobial skin cleanser for antiseptic handwashing on the ward, pre-operative surgical hand disinfection, and patient pre- and post-operative skin antiseptis.

Read the package leaflet before use, which is given when the label is peeled back.

Excipients: poloxamer 237, isopropyl alcohol, lauryl dimethyl amine oxide, glycerol, macrogol 7 glycerol cocoate, gluconolactone, citrus perfume, ponceau 4R (E124), sodium hydroxide and purified water.

Precautions: Avoid contact with eyes, brain, meninges and middle ear. Use with care in newborn babies, especially those born prematurely. Hibiscrub may cause chemical skin burns. Keep out of the sight and reach of children.

FOR EXTERNAL USE ONLY

Do not store above 25°C. Store in the original package.

Regent Medical Limited
Medlock Street, Oldham, Lancashire, OL1 3HS, UK

PL 22099/0001

Trade Mark of Regent Medical Ltd.



KSABF

5 060097 930760 >

PEEL HERE BUT DO NOT REMOVE

Page 5 - 91mm

Page 6 - 91mm

Front Cover - 99mm