


250 mm





250 mm

| you have been to d you have an intolerance to some sugars because this medicine also contains factose monohydrate (15.82mg per tablet). When administering this product to incontinent adults, pads should be changed more frequently to prevent extended skin contact with faeces. It is recommended to use changes in diet and bulk forming medicines to control constitution before using this medicine. Taking other medicines You should consult your doctor or pharmacist for advice if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. 3. How to take Senokot Max Strength Tablets 12 Years Plus New users should start with the lowest dose and increase it to the maximum if necessary. Once regularity has been regained dosage should be reduced and stopped. Adults, the elderly and children 12 years and over: swallow one tablet at night. Do not exceed the stated dose Duration of use | 4. Possible side effects Like all medicines, this medicine can cause side effects, although not everybody gets them. If any of the following or other side effects occur, stop taking this medicine and consult a doctor or pharmacist: - abdominal pain, spasms and diarrhoea, in particular these effects may be experienced by patients with irritable colon. In these cases, a reduction in dose may be neccessary. - hypersensitivity (allergic) reactions, hives (skin bumps), skin itchiness or local or more widespread, rash or asthma. - can make you more likely to catch infections. - chronic use may lead to albuminuria and haematuria which is albumin or red blood cells in the urine. - yellow or red-brown discoloration of urine may occur, which is harmless. - chronic use may cause pigmentation of the gastroinestinal tract which usually rededes when the product is no longer taken. - low potassium levels in the blood leading to muscle weakness and/or cramps - severe weight loss - changes in the shape of the finger and fingernail, muscular spasms and inflammatation of bones and joints The frequency is not known. |
|--|--|
| Normally it is sufficient to take this medicinal product up to two to three times a week. Not to be used for more than one week. Tell a doctor, pharmasist or nurse if: the symptoms worsen, or persist during the use of medicinal product. there is no bowel movement within 3 days of use. you need to take laxatives every day if abdominal pain persists. Do not take more than the leaflet tells you. Overuse can be harmful. | Reporting of side effects If you get any side effects, talk to your doctor, pharmasist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. 5. How to store |
| If you take too many tablets: Consult your doctor take this leafet with you. It may be harmful to take too much of this medicine or take it for too long. This is because it may lead to: 'lazy bowel' where the muscle in the bowel becomes too relaxed. This means that bowel emptying happens less often. This can lead to long-term constipation. imbalance of fluids and salts in the body. This can affect the tightness of muscle such as those in the bowel. It can also affect the salts in the blood. low levels of potassium in the blood (called hypo-kalaemia). This can make you tired, dizzy, make your muscles weak and cause an uneven heart beat. dehydration making you thirsty, feel faint and giving you headaches. It can also mean that you | Store below 30°C. Store in the original padkage. Do not use after the expiry date shown on the pack (EXP month/year). Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to Ithrow away medicines you no longer use. These measures will help protect the environment. 6. Contents of the pack and other information What Senokot Max Strength Tablets 12 Years Plus contains Each tablet contains powdered Alexandrian Senna fruit (pods) (cassia senna L. (C. acutifolia Delile)) and Tinnevelly Senna fruit (pods) (Cassia angustifolia Vahl) equivalent to 15mg hydroxyanthracene glycosides, calculated as sennoside B. Also contains: calcium phosphate, maize starch, magnesium stearate and lactose monohydrate. |
| cannot pass enough urine. If you miss a doset If you miss a dose, take your next dose at the usual time. Do not take two doses at the same time to make up for the missed dose. Always consult your doctor if you need further advice. | What your medicine looks like and the contents of the pack Senokot Max Strength Tablets 12 years Plus are available in pack sizes of 24 or 48 tablets. Not all pack sizes may be marketed. PL Holder and Manufacturer: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS IPL 00063/0770 Leafet last revised July 2020 Senokot is a trademark. 3159288 |

CUSTOMER INFO:
Minimum Point Size = 9.00pt

220

210

170

130

_0

-9

-6