Read these instructions carefully before taking this product—if you have any questions, ask your doctor or pharmacist. Pepto-Bismol oral suspension contains the active ingredient bismuth subsalicylate. It works by forming a soothing and protective coating of your stomach and intestine.

DO NOT USE IF YOU ARE: • under 16 years due to a possible reaction between salicylates and Reye's syndrome (a very rare but very serious disease) • allergic to salicylates including aspirin or any of the other ingredients • taking other salicylates including aspirin pregnant or breast-feeding.

### ASK A DOCTOR BEFORE USE IF YOU ARE TAKING MEDICINES FOR

- thinning the blood (anticoagulants) diabetes gout
- or tetracycline antibiotics.

## DIRECTIONS FOR USE: Adults and children over 16 years only:

Shake the bottle well before use. For oral use only. • Pour 30ml (one dose) into the cup provided and swallow. • Repeat the dose every 30-60 minutes if needed. • Do not take more than 8 doses in 24hrs • Do not exceed the stated dose or use after the expiry date. • Only use as directed. Use for longer than 2 days, or in higher doses than recommended has been associated with an increase in side effects. • If you have diarrhoea, it is important to prevent or treat dehydration by drinking plenty of fluids or taking oral rehydration products, especially if you are frail or elderly.





# Fast relief of:

- Heartburn
- Indigestion
- Nausea
- **▼Upset Stomach** Controls:
- Diarrhoea

## PROTECTIVE COATING ACTION

Oral Suspension 480ml

## Spare Out Location (To Contain Varnish and Color)

Lot & Exp ↑
P PL 00129/0358
Text revised 10/12



POSSIBLE SIDE EFFECTS: This product may cause a temporary and harmless darkening of the stool and/or tongue.

Warning this product contains amaranth (colouring) which may cause allergic type reactions including asthma.

STOP USING and SEE YOUR DOCTOR IF: • Symptoms are severe or persist for more than 2 days • you have diarrhoea accompanied by fever

● you experience ringing in the ears ● you notice anything unusual or have unexpected side effects • you accidentally take too much.

STORE: Below 25°C.

#### Keep out of the reach and sight of children.

Active is bismuth subsalicylate (526 mg in 30ml dose).

**Others:** aluminium magnesium silicate, methylcellulose, methyl salicylate, salicylic acid, sorbic acid, benzoic acid, saccharin sodium salicylate, amaranth and water.

Licence holder: Procter & Gamble, KT13 0XP

Release site: P&G, WN8 9NE, UK



XXXXXXX