

Patient Information Leaflet

Bonjela Complete Plus is a gel which once applied in the mouth forms a fine and soft film. This film is intended to:

- Protect the wound from external aggressions (food, etc)
- Soothe pain
- Aid healing

Indication:

Treatment of minor mouth ulcers exclusively (1 to 2 mouth ulcers < 1cm per fit) as well as small wounds of the oral cavity.

Contra-indication:

- Do not use in children under 16 years of age.
 This is because there is a possible association between the derivatives of salicylic acid and Reye's syndrome when given to children. Reye's syndrome is a very rare disease which affects the brain and liver and can be fatal.
- · Do not use in case of allergy to derivatives of salicylic acid.
- Do not use on major mouth ulcers (bigger than 1cm in diameter) or herpes-like ulcers (many tiny ulcers)
- Do not apply on infected wounds, burns or deep wounds (bleeding wounds).

Precautions for use:

- To be applied exclusively on the oral mucous membrane.
- May cause a temporary discomfort sensation and some tingling
- If the ulcer persist after 5 days of treatment, or if side effects occur speak to your doctor or pharmacist.

Directions for use:

- Wipe off the excess liquid from one side of the spatula against the rim of the bottle neck. Apply the product directly on the wound with the other side of the spatula in order to cover the area to treat,
- Allow to dry for about ten seconds before closing your mouth. A visible film will form over the ulcer. If the ulcer is not covered entirely, renew the application.
- Apply bonjela Complete Plus as often as necessary up to 4 times a day, preferably before meals, until healing (3 to 5 days).
- Never scratch off the film once formed, it will disappear by itself. So long as the film remains visible, it is not necessary to proceed to a new application.
- After use, replace the spatula into the bottle and close the bottle tightly. The composition of the gel guarantees its hygiene, it is therefore not necessary to clean the spatula after use.
- Seal tightly after use.
- The product should be used within 6 months after opening.

Ingredients:

Carboxylic acids, mineral acid, cellulose derivative, ethanol (anhydrous), purified water, orange flavour.

Storage:

- · Keep out of the reach of children.
- Do not use after the best-before date mentioned on the packaging.
- Keep the bottle tightly closed in a cool place
- Keep away from sources of ignition No smoking

Date of issue CE marking: 2010

Distributor: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS. Filmogel Technology is a technology licensed from Laboratories Urgo, France. bonjela is a trademark.

Manufacturer: S.D.P.P.A Route de Saulon. 21220 Gevrey-Chambertin, France



