



Quinoderm[®] cream

PLEASE READ THESE INSTRUCTIONS BEFORE USE.

Active Constituents:

BENZOYL PEROXIDE BP	10.0%
POTASSIUM HYDROXYQUINOLINE SULPHATE BP	0.5%

Other Constituents:

Lactic Acid BP, White Soft Paraffin BP, Edetic Acid BP, Sodium Acid Phosphate BP, Maize Starch BP, Cetyl Stearyl Alcohol, Sodium Cetyl Stearyl Sulphate, PEG 40 Castor Oil, Purified Water BP

Presentation

Quinoderm Cream is a creamy white vanishing cream and is available in tubes of 25g and 50g.

Mode of Action

Quinoderm Cream provides the peeling action of benzoyl peroxide and the antibacterial properties of potassium hydroxyquinoline sulphate.

Product Licence Holder

Ferndale Pharmaceuticals, 12 York Place, Leeds LS1 2DS, United Kingdom.

Manufacturer

Adams Healthcare (A division of Ecolab), Lotherton Way, Garforth, Leeds LS25 2JY, United Kingdom.

Indications

Acne vulgaris (common acne), acneform eruptions (acne like rash) and folliculitis (mild infection of the hair follicles).

When you should not use Quinoderm Cream

If in doubt about your skin problems discuss this with your doctor first. Patients with rosacea should not use Quinoderm Cream. Neither should those patients who have previously experienced an allergic reaction, such as a rash, itching or irritation, to any of the ingredients.

Precautions:

1. Quinoderm Cream should not be allowed to come into contact with the mouth or the eyes.

2. In a few isolated cases overreaction to Quinoderm Cream may occur. To minimise this possibility, select a small area of skin behind the ear, apply the cream and leave for twelve hours. If severe irritation or pronounced redness occurs do not proceed. If there is no reaction, treatment should be carried out as directed below.

3. Care should be taken to avoid contact with dyed fabrics as this preparation may adversely affect dye fastness.

Warnings:

In the unlikely event of an allergic reaction occurring, face oedema (swelling due to accumulation of fluid in the tissue spaces) may be a consequence. Under these circumstances consult your Doctor immediately.

Quinoderm Cream should not be used at the same time as other topical agents due to possible chemical interaction.

FOR EXTERNAL USE ONLY

Directions

IT IS IMPORTANT TO FOLLOW THESE INSTRUCTIONS CAREFULLY.

Adults, Children and the Elderly

Method of Application

Spread thinly over ALL the affected areas and gently massage until no trace of the cream remains. Apply two or three times daily or as directed by the physician (NB - It is important that the whole of the affected area be treated, not just individual spots or blemishes).

Side Effects

If your symptoms persist or if your condition worsens or if irritation, itch or rash occurs, stop treatment and consult your doctor or pharmacist. It is common for some slight redness of the skin to occur during the early weeks of treatment. This should gradually disappear as your skin becomes accustomed to the treatment. If any other undesirable effect occurs this should be reported to your doctor or pharmacist.

Do not use after expiry date:
refer to product labelling.

Store in dry conditions between 5°C and 30°C

Date of last review: April 2004

P2707-4